

Convex hulls of spheres and convex hulls of convex polytopes lying on parallel hyperplanes

Menelaos I. Karavelas^{†,‡} Eleni Tzanaki^{†,‡}

[†]*Department of Applied Mathematics,*

University of Crete

GR-714 09 Heraklion, Greece

`{mkaravel, etzanaki}@tem.uoc.gr`

[‡]*Institute of Applied and Computational Mathematics,*

Foundation for Research and Technology - Hellas,

P.O. Box 1385, GR-711 10 Heraklion, Greece

June 14, 2011

Abstract

Given a set Σ of spheres in $\mathbb{E}^d$, with $d \geq 3$ and d odd, having a fixed number of m distinct radii $\rho_1, \rho_2, \dots, \rho_m$, we show that the worst-case combinatorial complexity of the convex hull $CH_d(\Sigma)$ of Σ is $\Theta(\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor})$, where n_i is the number of spheres in Σ with radius ρ_i . Our bound refines the worst-case upper and lower bounds on the worst-case combinatorial complexity of $CH_d(\Sigma)$ for all odd $d \geq 3$.

To prove the lower bound, we construct a set of $\Theta(n_1 + n_2)$ spheres in $\mathbb{E}^d$, with $d \geq 3$ odd, where n_i spheres have radius ρ_i , $i = 1, 2$, and $\rho_2 \neq \rho_1$, such that their convex hull has combinatorial complexity $\Omega(n_1 n_2^{\lfloor \frac{d}{2} \rfloor} + n_2 n_1^{\lfloor \frac{d}{2} \rfloor})$. Our construction is then generalized to the case where the spheres have $m \geq 3$ distinct radii.

For the upper bound, we reduce the sphere convex hull problem to the problem of computing the worst-case combinatorial complexity of the convex hull of a set of m d -dimensional convex polytopes lying on m parallel hyperplanes in $\mathbb{E}^{d+1}$, where $d \geq 3$ odd, a problem which is of independent interest. More precisely, we show that the worst-case combinatorial complexity of the convex hull of a set $\{\mathcal{P}_1, \mathcal{P}_2, \dots, \mathcal{P}_m\}$ of m d -dimensional convex polytopes lying on m parallel hyperplanes of $\mathbb{E}^{d+1}$ is $O(\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor})$, where n_i is the number of vertices of $\mathcal{P}_i$. This bound is an improvement over the worst-case bound on the combinatorial complexity of the convex hull of a point set where we impose no restriction on the points' configuration; using the lower bound construction for the sphere convex hull problem, it is also shown to be tight for all odd $d \geq 3$.

Finally: (1) we briefly discuss how to compute convex hulls of spheres with a fixed number of distinct radii, or convex hulls of a fixed number of polytopes lying on parallel hyperplanes; (2) we show how our tight bounds for the parallel polytope convex hull problem, yield tight bounds on the combinatorial complexity of the Minkowski sum of two convex polytopes in $\mathbb{E}^d$; and (3) we state some open problems and directions for future work.

Key words: high-dimensional geometry, discrete geometry, combinatorial geometry, combinatorial complexity, convex hull, Minkowski sum, spheres, convex polytopes, parallel hyperplanes

2010 MSC: 68U05 (Primary), 52B05, 52B11, 52C45 (Secondary)

1 Introduction and results

Let Σ be a set of n spheres in $\mathbb{E}^d$, $d \geq 2$, where the dimension d is fixed. We call Π a *supporting* hyperplane of Σ if it has non-empty intersection with Σ and Σ is contained in one of the two closed halfspaces bounded by Π . We call H a *supporting halfspace* of the set Σ if it contains all spheres in Σ and is limited by a supporting hyperplane Π of Σ . The intersection of all supporting halfspaces of Σ is called the convex hull $CH_d(\Sigma)$ of Σ . The definition of convex hulls detailed above is applicable not only to spheres, but also to any finite set of compact geometric objects in $\mathbb{E}^d$. In the case of points, i.e., if we have a set P of n points in $\mathbb{E}^d$, the worst-case combinatorial complexity¹ of $CH_d(P)$ is known to be $\Theta(n^{\lfloor \frac{d}{2} \rfloor})$. Moreover, there exist worst-case optimal algorithms for constructing $CH_d(P)$ that run in $O(n^{\lfloor \frac{d}{2} \rfloor} + n \log n)$ time, e.g., see [16, 25, 2, 24, 13]. Since the complexity of $CH_d(P)$ may vary from $O(1)$ to $\Theta(n^{\lfloor \frac{d}{2} \rfloor})$, a lot of work has been devoted to the design of output-sensitive algorithms for constructing $CH_d(P)$, i.e., algorithms the running time of which depends on the size of the output convex hull $CH_d(P)$, e.g., see [12, 18, 27, 19, 21, 14, 1, 9, 10, 11]. For a nice overview of the various algorithms for computing the convex hull of points sets, the interested reader may refer to the paper by Erickson [15], while Avis, Bremner and Seidel [4] have a very nice discussion about the effectiveness of output-sensitive convex hull algorithms for point sets.

Results about the convex hull of non-linear objects are very limited. Aurenhammer [3] showed that the worst-case complexity of the power diagram of a set of n spheres in $\mathbb{E}^d$, $d \geq 2$, is $O(n^{\lceil \frac{d}{2} \rceil})$, which also implies the same upper bound for the worst-case complexity for the convex hull of the sphere set. Rappaport [26] devised an $O(n \log n)$ algorithm for computing the convex hull of a set of discs on the plane, which is worst-case optimal. Boissonnat *et al.* [6] give an $O(n^{\lceil \frac{d}{2} \rceil} + n \log n)$ algorithm for computing the convex hull of a set of n spheres in $\mathbb{E}^d$, $d \geq 2$, which is worst-case optimal in three and even dimensions, since they also show that the worst-case complexity of the convex hull of n spheres in $\mathbb{E}^3$ is $\Theta(n^2)$. Finally, their results hold true for the case of homothetic convex objects. Boissonnat and Karavelas [7] settled a conjecture in [6]: they proved that the worst-case complexity of a set of n spheres in $\mathbb{E}^d$, $d \geq 2$, is $\Theta(n^{\lceil \frac{d}{2} \rceil})$, which also implied that the algorithm presented in [6] is optimal for all d . As far as output-sensitive algorithms are concerned, Boissonnat, C  r  zo and Duquesne [5] showed how to construct the convex hull of a set of n three-dimensional spheres in $O(nf)$ time, where f is the size of the output convex hull, while Nielsen and Yvinec [23] discuss optimal or almost optimal output-sensitive convex hull algorithms for planar convex objects.

In this paper we consider the problem of computing the complexity of the convex hull of a set of spheres, when the spheres have a fixed number of distinct radii. This problem has been posed by Boissonnat and Karavelas [7], and it is meaningful for odd dimensions only: in even dimensions the complexity of both the convex hull of n points and the convex hull of n spheres is $\Theta(n^{\lfloor \frac{d}{2} \rfloor}) = \Theta(n^{\lceil \frac{d}{2} \rceil})$, i.e., the two bounds match.

Consider a set of n spheres Σ in $\mathbb{E}^d$, where $d \geq 3$ and d odd, such that the spheres in Σ have a fixed number m of distinct radii $\rho_1, \rho_2, \dots, \rho_m$. Let n_i be the number of spheres in Σ with radius ρ_i . We say that ρ_λ *dominates* Σ if $n_\lambda = \Theta(n)$. We further say that Σ is *uniquely* (resp., *strongly*) *dominated*, if, for some λ , ρ_λ dominates Σ , and $n_i = o(n)$ (resp., $n_i = O(1)$), for all $i \neq \lambda$. Using this terminology, we can qualitatively express our results as follows. Firstly, if Σ is strongly dominated, then, from the combinatorial complexity point of view, $CH_d(\Sigma)$ behaves as if we had a set of points, or equivalently a set of spheres with the same radius. If, however, Σ is dominated by at least two radii, $CH_d(\Sigma)$ behaves as in the generic case, where we impose no restriction on the number of distinct radii in Σ . Finally, if Σ is uniquely dominated (but not strongly dominated), the complexity

¹In the rest of the paper, and unless otherwise stated, we use the term “complexity” to refer to “combinatorial complexity”.

of $CH_d(\Sigma)$ stands in-between the two extremes above: the complexity of $CH_d(\Sigma)$ is asymptotically larger than the points' case (or when we have spheres with the same radius), and asymptotically smaller than the generic case, where we impose no restriction on the number of distinct radii in Σ . From the quantitative point of view, our results refine the results in [7] for any odd dimension $d \geq 3$ as follows. We prove that the worst-case complexity of $CH_d(\Sigma)$ is $\Theta(\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor})$. This tight bound constitutes an improvement over the generic worst-case complexity of $CH_d(\Sigma)$ if Σ is uniquely dominated, since in this case the complexity of $CH_d(\Sigma)$ is $o(n^{\lfloor \frac{d}{2} \rfloor})$. On the other hand, it matches the generic worst-case complexity of $CH_d(\Sigma)$, if Σ is dominated by at least two radii: in this case the worst-case complexity of $CH_d(\Sigma)$ becomes $\Theta(n^{\lfloor \frac{d}{2} \rfloor})$. Finally, if Σ is strongly dominated, the complexity of $CH_d(\Sigma)$ is $\Theta(n^{\lfloor \frac{d}{2} \rfloor})$, i.e., it matches the worst-case complexity of convex hulls of point sets (or sets of spheres where all spheres have the same radius).

To establish the lower bound for the complexity of $CH_d(\Sigma)$, we construct a set Σ of $\Theta(n_1 + n_2)$ spheres in $\mathbb{E}^d$, for any odd $d \geq 3$, where n_1 spheres have radius ρ_1 and n_2 spheres have radius $\rho_2 \neq \rho_1$, such that worst-case complexity of $CH_d(\Sigma)$ is $\Omega(n_1 n_2^{\lfloor \frac{d}{2} \rfloor} + n_2 n_1^{\lfloor \frac{d}{2} \rfloor})$. This construction is then generalized to sets of spheres having a fixed number of $m \geq 3$ distinct radii. More precisely, we construct a set Σ of $n = \sum_{i=1}^m n_i$ spheres, where n_i spheres have radius ρ_i , with the ρ_i 's being pairwise distinct, such that the worst-case complexity of $CH_d(\Sigma)$ is $\Omega(\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor})$.

To prove our upper bound we use a lifting map, introduced in [6], that lifts spheres $\sigma_i = (c_i, r_i)$ in $\mathbb{E}^d$ to points $p_i = (c_i, r_i)$ in $\mathbb{E}^{d+1}$. The convex hull $CH_d(\Sigma)$ is then the intersection of the hyperplane $\{x_{d+1} = 0\}$ with the Minkowski sum of the convex hull $CH_{d+1}(P)$ and the hypercone λ_0 , where P is the point set $\{p_1, p_2, \dots, p_n\}$ in $\mathbb{E}^{d+1}$, and λ_0 is the lower half hypercone with arbitrary apex, vertical axis and angle at the apex equal to $\frac{\pi}{4}$. When the spheres in Σ have a fixed number m of distinct radii, the points of P lie on m hyperplanes parallel to the hyperplane $\{x_{d+1} = 0\}$. In this setting, computing the complexity of $CH_d(\Sigma)$ amounts to computing the complexity of the convex hull of m convex polytopes lying on m parallel hyperplanes of $\mathbb{E}^{d+1}$. This observation gives rise to the second major result in this paper, which is of independent interest, and gives as corollary a tight bound on the worst-case complexity of the Minkowski sum of two convex d -polytopes. Given a set $\mathcal{P} = \{\mathcal{P}_1, \mathcal{P}_2, \dots, \mathcal{P}_m\}$ of m convex d -polytopes in $\mathbb{E}^{d+1}$, with $d \geq 3$ and d odd, we show that the worst-case complexity of the convex hull $CH_{d+1}(\mathcal{P})$ is $\Theta(\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor})$, where n_i is the number of vertices of $\mathcal{P}_i$. Our upper bound proof is by induction on the number m of parallel hyperplanes. The lower bound follows from the lower bound on the complexity of the convex hull of spheres having m distinct radii. Our bound constitutes an improvement over the worst-case complexity of convex hulls of points sets, if a single polytope of $\mathcal{P}$ has $\Theta(n)$ vertices, whereas all other polytopes have $o(n)$ vertices, where n is the total number of vertices of all m polytopes, while it matches the worst-case complexity of convex hulls of points sets if at least two polytopes have $\Theta(n)$ vertices.

The rest of our paper is structured as follows: In Section 2 we detail our inductive proof of the upper bound on the worst-case complexity of the convex hull of convex polytopes lying on parallel hyperplanes, while in Section 3 we discuss how to compute this convex hull. In Section 4 we prove our upper bound on the worst-case complexity of the convex hull of a set of spheres. Next we present our lower bound construction for any odd $d \geq 3$ in two steps: first for sphere sets with two distinct radii and then for sphere sets with $m \geq 3$ distinct radii. We end the section by discussing how this lower bound yields a tight lower bound for the problem of the previous section. In Section 5 we explain how to modify the algorithm by Boissonnat *et al.* [6] so as to almost optimally compute the convex hull of a set of spheres with a fixed number of distinct radii. Finally, in Section 6 we summarize our results, we explain how our results yield tight bounds for the complexity of the

Minkowski sum of two convex polytopes, and state some open problems.

2 Convex hulls of convex polytopes lying on parallel hyperplanes

A *convex polytope*, or simply *polytope*, $\mathcal{P}$ in $\mathbb{E}^d$ is the convex hull of a finite set of points P in $\mathbb{E}^d$. A polytope $\mathcal{P}$ can equivalently be described as the intersection of all the closed halfspaces containing P . A *face* of $\mathcal{P}$ is an intersection of $\mathcal{P}$ with hyperplanes for which the polytope is contained in one of the two closed halfspaces determined by the hyperplane. The dimension of a face of $\mathcal{P}$ is the dimension of its affine hull. A k -face of $\mathcal{P}$ is a k -dimensional face of $\mathcal{P}$. We consider the polytope itself as a trivial d -dimensional face; all the other faces are called *proper* faces. We will use the term *d-polytope* to refer to a polytope the trivial face of which is d -dimensional. For a d -polytope $\mathcal{P}$, the 0-faces of $\mathcal{P}$ are its *vertices*, the 1-faces of $\mathcal{P}$ are its *edges*, the $(d-2)$ -faces of $\mathcal{P}$ are called *ridges*, while the $(d-1)$ -faces are called *facets*. For $0 \leq k \leq d$ we denote by $f_k(\mathcal{P})$ the number of k -faces of $\mathcal{P}$. Note that every k -face F of $\mathcal{P}$ is also a k -polytope whose faces are all the faces of $\mathcal{P}$ contained in F . A k -simplex in $\mathbb{E}^d$, $k \leq d$, is the convex hull of any $k+1$ affinely independent points in $\mathbb{E}^d$. A polytope is called *simplicial* if all its proper faces are simplices. Equivalently, $\mathcal{P}$ is simplicial if for every vertex v of $\mathcal{P}$ and every face $F \in \mathcal{P}$, v does not belong to the affine hull of the vertices in $F \setminus \{v\}$.

A *polytopal complex* $\mathcal{C}$ is a finite collection of polytopes in $\mathbb{E}^d$ such that (i) $\emptyset \in \mathcal{C}$, (ii) if $\mathcal{P} \in \mathcal{C}$ then all the faces of $\mathcal{P}$ are also in $\mathcal{C}$ and (iii) the intersection $\mathcal{P} \cap \mathcal{Q}$ for two polytopes in $\mathcal{C}$ is a face of both $\mathcal{P}$ and $\mathcal{Q}$. The dimension $\dim(\mathcal{C})$ of $\mathcal{C}$ is the largest dimension of a polytope in $\mathcal{C}$. A polytopal complex is called *pure* if all its maximal (with respect to inclusion) faces have the same dimension. In this case the maximal faces are called the *facets* of $\mathcal{C}$. We will use the term *d-complex* to refer to a pure polytopal complex whose facets are d -dimensional. A polytopal complex is simplicial if all its faces are simplices. Finally, a polytopal complex $\mathcal{C}'$ is called a subcomplex of a polytopal complex $\mathcal{C}$ if all faces of $\mathcal{C}'$ are also faces of $\mathcal{C}$.

One important class of polytopal complexes arise from polytopes. More precisely, a d -polytope $\mathcal{P}$, together with all its faces and the empty set, form a polytopal d -complex, denoted by $\mathcal{C}(\mathcal{P})$. The only maximal face of $\mathcal{C}(\mathcal{P})$, which is clearly the only facet of $\mathcal{C}(\mathcal{P})$, is the polytope $\mathcal{P}$ itself. Moreover, all proper faces of $\mathcal{P}$ form a pure polytopal complex, called the *boundary complex* $\mathcal{C}(\partial\mathcal{P})$. The facets of $\mathcal{C}(\partial\mathcal{P})$ are just the facets of $\mathcal{P}$, and its dimension is $\dim(\mathcal{P}) - 1 = d - 1$. Given a polytope $\mathcal{P}$ and a vertex v of $\mathcal{P}$, the *star* of v is the polytopal complex of all faces of $\mathcal{P}$ that contain v , and their faces. The *link* of v is the subcomplex of the star of v consisting of all the faces of the star of v that do not contain v .

The f -vector $(f_{-1}(\mathcal{P}), f_0(\mathcal{P}), \dots, f_{d-1}(\mathcal{P}))$ of a simplicial d -polytope $\mathcal{P}$ is defined as the $(d+1)$ -dimensional vector consisting of the number $f_k(\mathcal{P})$ of k -faces of $\mathcal{P}$, $-1 \leq k \leq d$, where $f_{-1}(\mathcal{P}) = 1$ refers to the empty set. The h -vector $(h_0(\mathcal{P}), h_1(\mathcal{P}), \dots, h_d(\mathcal{P}))$ of a simplicial d -polytope $\mathcal{P}$ is defined as the $(d+1)$ -dimensional vector, where $h_k(\mathcal{P}) := \sum_{i=0}^k (-1)^{k-i} \binom{d-i}{d-k} f_{i-1}(\mathcal{P})$, $0 \leq k \leq d$. The number $h_k(\mathcal{P})$ counts the number of facets of $\mathcal{P}$ in a *shelling* of $\mathcal{P}$, whose *restriction* has size k ; this number is independent of the particular shelling chosen (cf. [28, Theorem 8.19]). It is easy to verify from the defining equations of the $h_k(\mathcal{P})$'s that the elements of the f -vector determine the elements of the h -vector and vice versa. Moreover, the elements of the f -vector (or, equivalently, the h -vector) are not linearly independent; they satisfy the so called *Dehn-Sommerville equations*, which can be written in a very concise form as: $h_k(\mathcal{P}) = h_{d-k}(\mathcal{P})$, $0 \leq k \leq d$. An important implication of the existence of the Dehn-Sommerville equations is that if we know the face numbers $f_k(\mathcal{P})$ for all $0 \leq k \leq \lfloor \frac{d}{2} \rfloor - 1$, we can determine the remaining face numbers $f_k(\mathcal{P})$ for all $\lfloor \frac{d}{2} \rfloor \leq k \leq d-1$.

In what follows we recall some facts from [17, Section 5.2] that will be of use to us later. Let $\mathcal{P}$

be a d -polytope in $\mathbb{E}^d$, F a facet of $\mathcal{P}$, and H the supporting hyperplane of F (with respect to $\mathcal{P}$). For an arbitrary point p in $\mathbb{E}^d$, we say that p is *beyond* (resp., *beneath*) the facet F of $\mathcal{P}$, if p lies in the open halfspace of H that does not contain $\mathcal{P}$ (resp., contains the interior of $\mathcal{P}$). Furthermore, we say that an arbitrary point v' is *beyond* the vertex v of $\mathcal{P}$ if for every facet F of $\mathcal{P}$ containing v , v' is beyond F , while for every facet F of $\mathcal{P}$ not containing v , v' is beneath F . The vertices of the polytope $\mathcal{P}' = CH_d((P \setminus \{v\}) \cup \{v'\})$ are the same with those of $\mathcal{P}$, except for v which has been replaced by v' . In this case we say that $\mathcal{P}'$ is obtained from $\mathcal{P}$ by *pulling* v to v' . The vertex v' of $\mathcal{P}'$ does not belong to the affine hull of the vertices in $F' \setminus \{v'\}$ for every face F' of $\mathcal{P}'$. The following result is well-known.

Theorem 1 ([20, 22]). *Let $\mathcal{P}$ be a d -polytope.*

- (i) *The d -polytope $\mathcal{P}'$ we obtain by pulling a vertex of $\mathcal{P}$ has the same number of vertices with $\mathcal{P}$, and $f_k(\mathcal{P}) \leq f_k(\mathcal{P}')$ for all $1 \leq k \leq d-1$.*
- (ii) *The d -polytope $\mathcal{P}'$ we obtain by successively pulling each of the vertices of $\mathcal{P}$ is simplicial, has the same number of vertices with $\mathcal{P}$, and $f_k(\mathcal{P}) \leq f_k(\mathcal{P}')$ for all $1 \leq k \leq d-1$.*

In the rest of the paper, when we refer to parallel hyperplanes we assume that they have the same unit normal vector, i.e., they have the same orientation. Moreover, if two hyperplanes Π and Π' are parallel, we say that Π' is *above* Π if Π' lies in the positive open halfspace delimited by Π .

Let $\mathcal{S} = \{\mathcal{P}_1, \mathcal{P}_2, \dots, \mathcal{P}_m\}$ be a set of m d -polytopes lying on m parallel hyperplanes $\Pi_1, \Pi_2, \dots, \Pi_m$ of $\mathbb{E}^{d+1}$, respectively. Throughout this section we assume that $m \geq 2$ is fixed, and that Π_j is above Π_i for all $j > i$. We denote by P_i the set of vertices of $\mathcal{P}_i$, by n_i the cardinality of P_i , and by P the union $P = P_1 \cup P_2 \cup \dots \cup P_m$. Let $\mathcal{P} = CH_{d+1}(P)$; note that, for each i , not all vertices in P_i are necessarily vertices of $\mathcal{P}$. Furthermore, among the polytopes in $\mathcal{S}$, only $\mathcal{P}_1$ and $\mathcal{P}_m$ are faces of $\mathcal{P}$.

The theorem that follows is the adaptation of Theorem 1 in the context of m parallel polytopes. Again, we want to perturb the points in P so that $\mathcal{P}'$ (the polytope we obtain after perturbing the points in P) is simplicial with $f_k(\mathcal{P}) \leq f_k(\mathcal{P}')$ for $k \geq 1$, but we want to retain the property that the points lying on a hyperplane Π_i , if perturbed, are replaced by points that lie on the same hyperplane. This is almost possible. More precisely, all the faces of the polytope $\mathcal{P}'$ are simplicial, with the possible exception of $\mathcal{P}'_1 = \mathcal{P}' \cap \Pi_1$ and $\mathcal{P}'_m = \mathcal{P}' \cap \Pi_m$.

Lemma 2. *Let $\mathcal{S} = \{\mathcal{P}_1, \mathcal{P}_2, \dots, \mathcal{P}_m\}$ be a set of $m \geq 2$ d -polytopes lying on m parallel hyperplanes $\Pi_1, \Pi_2, \dots, \Pi_m$ of $\mathbb{E}^{d+1}$, respectively, where Π_j is above Π_i for all $j > i$. Let P_i be the vertex set of $\mathcal{P}_i$, $1 \leq i \leq m$, $P = P_1 \cup P_2 \dots \cup P_m$, and $\mathcal{P} = CH_{d+1}(P)$. The points in P can be perturbed in such a way that:*

- (i) *the points of P in each hyperplane Π_i remain in Π_i , $1 \leq i \leq m$,*
- (ii) *all the faces of $\mathcal{P}'$, except possibly the facets $\mathcal{P}'_1$ and $\mathcal{P}'_m$, are simplices, and,*
- (iii) *$f_k(\mathcal{P}) \leq f_k(\mathcal{P}')$ for all $1 \leq k \leq d$,*

where $\mathcal{P}'$ is the polytope we obtain after having perturbed the vertices of $\mathcal{P}$ in P .

Proof. We construct $\mathcal{P}'$ in three steps. Firstly, we properly perturb the points in Π_1 so that $\mathcal{P}'_1$ is simplicial, then we do the same for the points in Π_m , and, finally, we pull every vertex of $\mathcal{P}$ in Π_i for $2 \leq i \leq m-1$.

Let $v \in \Pi_1$ and choose any v' in Π_1 beyond all the facets of $\mathcal{P}$ other than $\mathcal{P}_1$ that contain v . If we focus our attention on the hyperplane Π_1 and on the polytope $\mathcal{P}_1 \in \Pi_1$ we have that v is a vertex of $\mathcal{P}_1$ and v' is some point beyond all the facets of $\mathcal{P}_1$ containing v . We consider the polytope $\mathcal{P}' = CH_{d+1}((P \setminus \{v\}) \cup \{v'\})$. In view of [17, Theorems 5.1.1 & 5.1.2] the k -faces of $\mathcal{P}'$ are either:

- (i) k -faces of $\mathcal{P}'_1$, or

- (ii) k -faces of $\mathcal{P}$ not in Π_1 that do not contain v , or
- (iii) faces of the form $CH_{d+1}(\{v'\} \cup G_{k-1})$, where G_{k-1} is a $(k-1)$ -face not containing v of a facet $F \notin \Pi_1$ of $\mathcal{P}$ containing v .

The number of k -faces of $\mathcal{P}'$ does not change due to Case (ii). The number of k -faces of $\mathcal{P}'$ may increase in Case (iii) if G_{k-1} is contained in a non-simplicial k -face of $\mathcal{P}$, otherwise it does not increase. Finally, Case (i) is a consequence of Theorem 1(i).

We set $\mathcal{P} := \mathcal{P}'$ and we repeat the above procedure for every point $v \in P_1$. After having perturbed all points in P_1 we obtain a polytope $\mathcal{P}'$ such that $\mathcal{P}'_1$ is a simplicial polytope. Moreover, we have the additional properties that $f_k(\mathcal{P}) \leq f_k(\mathcal{P}')$, for $k \geq 1$, and that for every $v' \in \Pi_1$ and for each face $F' \in \mathcal{P}'$ not lying in Π_1 , the point v' does not lie in the affine hull of the vertices in $F' \setminus \{v'\}$. We repeat exactly the same procedure on the points of Π_m . Again, we denote by $\mathcal{P}$ the polytope we obtain after having perturbed the points in P_1 and P_m .

It is left to pull the points in the remaining hyperplanes; in fact, we need only perturb the points in P that are vertices of $\mathcal{P}$. Consider a vertex v of $\mathcal{P}$, such that $v \in P_i$ for some $2 \leq i \leq m-1$. Let v' be any point in Π_i that lies beyond all the facets of $\mathcal{P}$ containing v . The choice of such a v' is possible since Π_i is a hyperplane containing v but not a supporting hyperplane of $\mathcal{P}$. The polytope $\mathcal{P}' = CH_{d+1}((P \setminus \{v\}) \cup \{v'\})$ is the one we obtain by pulling the vertex v to v' and thus from Theorem 1(i) we have $f_k(\mathcal{P}) \leq f_k(\mathcal{P}')$ for all $k \geq 1$. We continue the same procedure for every vertex v of $\mathcal{P}$ in Π_i , and $2 \leq i \leq m-1$. After having pulled all the vertices of $\mathcal{P}$ in Π_i , for $2 \leq i \leq m-1$, we obtain a polytope $\mathcal{P}'$ with the property that for every vertex v' of $\mathcal{P}'$ in Π_i , with $2 \leq i \leq m-1$, and for every face $F' \in \mathcal{P}'$, the point v' does not lie in the affine hull of the vertices in $F' \setminus \{v'\}$.

Summarizing all of the above, we deduce that, after having perturbed all the vertices of $\mathcal{P}$ in P , we get a polytope $\mathcal{P}'$ with the same number of vertices as $\mathcal{P}$, such that: (1) $f_k(\mathcal{P}) \leq f_k(\mathcal{P}')$ for all $k \geq 1$, (2) $\mathcal{P}'_1$ and $\mathcal{P}'_m$ are simplicial, and (3) for every vertex v' of $\mathcal{P}'$ and every face $F' \in \mathcal{P}'$, such that F' does not lie in Π_1 or Π_m , the vertex v' does not lie in the affine hull of the vertices in $F' \setminus \{v'\}$. The latter implies that all the faces of $\mathcal{P}'$, not in Π_1 or Π_m , are simplices. This completes our proof. $\square$

In view of Lemma 2, it suffices to restrict our attention to sets of polytopes $\mathcal{P}$, where $\mathcal{P}$ is simplicial with the possible exception of its two facets $\mathcal{P}_1$ and $\mathcal{P}_m$. Let $\tilde{\Pi}$ be any hyperplane between and parallel to the hyperplanes Π_{m-1} and Π_m and consider the intersection $\tilde{\mathcal{P}} := \mathcal{P} \cap \tilde{\Pi}$ (see Fig. 1). Let $\mathcal{F}$ be the set of faces of $\mathcal{P}$ having non-empty intersection with $\tilde{\Pi}$. Note that $\tilde{\mathcal{P}}$ is a d -polytope, which is, in general, non-simplicial, and whose proper non-trivial faces are intersections of the form $F \cap \tilde{\Pi}$ where $F \in \mathcal{F}$. Let $A = (\alpha_1, \alpha_2, \dots, \alpha_m)$ and $B = (\beta_1, \beta_2, \dots, \beta_m)$ be two vectors in $\mathbb{N}^m$. We say that $A \preceq B$ if $\alpha_i \leq \beta_i$ for all $1 \leq i \leq m$, and denote by $|A|$ the sum of the elements of A , i.e., $|A| = \sum_{i=1}^m \alpha_i$. The following lemma provides an upper bound on the number of k -faces of $\mathcal{F}$.

Lemma 3. *The number of k -faces of $\mathcal{F}$ is bounded from above as follows:*

$$f_k(\mathcal{F}) \leq \sum_{\substack{(0, \dots, 0, 1) \preceq A \preceq (k, \dots, k) \\ |A|=k+1}} \prod_{i=1}^m \binom{f_0(\mathcal{P}_i)}{\alpha_i}, \quad 1 \leq k \leq d,$$

where α_i is the i -th coordinate of the vector $A \in \mathbb{N}^m$.

Proof. According to Lemma 2, it suffices to consider the case where $\mathcal{P}$ is simplicial except possibly for its facets $\mathcal{P}_1$ and $\mathcal{P}_m$. In this context, a k -face $F \in \mathcal{F}$ is simplicial and it is defined by $k+1$ vertices of P , where at least one vertex comes from P_m , whereas the remaining k vertices are vertices




Figure 1: An example of four polytopes lying on four parallel hyperplanes of $\mathbb{E}^d$. The white vertices are not vertices of $\mathcal{P}$. The polytopes $\mathcal{P}_2$ and $\mathcal{P}_3$ are shown in black. All faces of $\mathcal{L}$ are shown in green (only the facet $\mathcal{P}_1$ is shown), whereas all faces of $\mathcal{P}_4$ are shown in red. The faces in blue are faces of $\mathcal{F}$, whereas the faces in brown are faces of $\tilde{\mathcal{P}} = \mathcal{P} \cap \tilde{\Pi}$.

of $P_1, P_2, \dots, P_{m-1}$. Let α_i be the number of vertices of F from P_i . Clearly, we have $0 \leq \alpha_i \leq k$ for $1 \leq i \leq m-1$, $1 \leq \alpha_m \leq k$, and $\alpha_1 + \alpha_2 + \dots + \alpha_m = k+1$. The maximum number of possible $(\alpha_i - 1)$ -faces of $\mathcal{P}_i$ is $\binom{f_0(\mathcal{P}_i)}{\alpha_i}$, which implies that the maximum possible number of k -faces of $\mathcal{F}$ is $\prod_{i=1}^m \binom{f_0(\mathcal{P}_i)}{\alpha_i}$. Summing over all possible values for the α_i 's we get the desired expression. $\square$

Using the bounds for the face numbers of $\mathcal{F}$ from Lemma 3, we arrive at the following lemma concerning bounds on the number of k -faces of $\mathcal{P}$, for small values of k .

Lemma 4. *Let $n_i = f_0(\mathcal{P}_i)$, $1 \leq i \leq m$. The following asymptotic bounds hold:*

$$f_k(\mathcal{P}) = O\left(\sum_{1 \leq i \neq j \leq m} n_i n_j^k + \sum_{i=1}^m n_i^{\min\{k+1, \lfloor \frac{d}{2} \rfloor\}}\right), \quad 0 \leq k \leq \lfloor \frac{d}{2} \rfloor.$$

Proof. In what follows k will be at most $\lfloor \frac{d}{2} \rfloor$, while we denote by k' the quantity $k' = \min\{k+1, \lfloor \frac{d}{2} \rfloor\}$. Let $T(m)$ be the worst-case complexity of $f_k(\mathcal{P})$. We are going to prove, by induction on m , that

for all $m \geq 1$, $T(m) \leq c(\sum_{1 \leq i \neq j \leq m} n_i n_j^k + \sum_{i=1}^m n_i^{k'})$, where c is some appropriately large constant that depends only on d .

The case $m = 1$ is trivial since the number of k -faces of a d -polytope with n_1 vertices is in $O(n_1^{k'})$. Let us now assume that $m \geq 2$ and that our statement holds for $m - 1$; we shall prove it for m . To this end, we consider a set of m d -polytopes $\mathcal{P} = \{\mathcal{P}_1, \mathcal{P}_2, \dots, \mathcal{P}_m\}$, lying on m parallel hyperplanes $\Pi_1, \dots, \Pi_m$ of $\mathbb{E}^{d+1}$, such that Π_j is above Π_i for all $j > i$. Let $n_i = f_0(\mathcal{P}_i)$, $1 \leq i \leq m$, and denote by $\mathcal{P}$ the convex hull $CH_{d+1}(\mathcal{P})$. Consider a hyperplane $\tilde{\Pi}$ parallel to Π_{m-1} and Π_m and between them, and let $\tilde{\mathcal{P}} = \mathcal{P} \cap \tilde{\Pi}$ (refer to Fig. 1). We denote by $\mathcal{L}$ the set of faces F of $\mathcal{P}$ such that $F \not\subset \mathcal{P}_m$ and $F \cap \tilde{\Pi} = \emptyset$, and by $\mathcal{F}$ the set of faces F of $\mathcal{P}$ with $F \cap \tilde{\Pi} \neq \emptyset$. Clearly, the set of faces of $\mathcal{P}$ is equal to the disjoint union of $\mathcal{L}$, $\mathcal{F}$ and the set of faces of $\mathcal{P}_m$; hence:

$$f_k(\mathcal{P}) = f_k(\mathcal{L}) + f_k(\mathcal{F}) + f_k(\mathcal{P}_m). \quad (1)$$

By the induction hypothesis we have that the number of k -faces of $CH_{d+1}(\mathcal{P} \setminus \{\mathcal{P}_m\})$ is at most $T(m - 1)$. Since the k -faces in $\mathcal{L}$ are k -faces of $CH_{d+1}(\mathcal{P} \setminus \{\mathcal{P}_m\})$, we have that $f_k(\mathcal{L})$ is at most $T(m - 1)$. On the other hand, by Lemma 3, we immediately get that $f_k(\mathcal{F}) = O(n_m^k \sum_{i=1}^{m-1} n_i + n_m \sum_{i=1}^{m-1} n_i^k)$. Finally, since $\mathcal{P}_m$ is a d -polytope, $f_k(\mathcal{P}_m) = O(n_m^{k'})$. Combining these bounds with eq. (1), we arrive at the following recurrence relation for $T(m)$:

$$T(m) \leq T(m - 1) + O(n_m^k \sum_{i=1}^{m-1} n_i + n_m \sum_{i=1}^{m-1} n_i^k) + O(n_m^{k'}).$$

It is straightforward to verify that $T(m)$ satisfies: $T(m) \leq c(\sum_{1 \leq i \neq j \leq m} n_i n_j^k + \sum_{i=1}^m n_i^{k'})$, for some appropriately large constant c (that depends only on d); this establishes our claim. $\square$

Exploiting the bounds from Lemma 4 for $0 \leq k \leq \lfloor \frac{d}{2} \rfloor$, and using the Dehn-Sommerville equations of an appropriately defined simplicial $(d + 1)$ -polytope containing all faces in $\mathcal{P}$, except for the facets $\mathcal{P}_1$ and $\mathcal{P}_m$, we derive asymptotic bounds on $f_k(\mathcal{P})$, for all $\lfloor \frac{d+1}{2} \rfloor \leq k \leq d$. Our results are summarized in the following lemma.

Lemma 5. *Let $n_i = f_0(\mathcal{P}_i)$, $1 \leq i \leq m$. The following asymptotic bounds hold:*

$$f_k(\mathcal{P}) = O\left(\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor}\right), \quad \lfloor \frac{d+1}{2} \rfloor \leq k \leq d.$$

Proof. Let y (resp., z) be a point below Π_1 (resp., above Π_m), such that the vertices of $\mathcal{P}_1$ (resp., $\mathcal{P}_m$) are the only vertices of $\mathcal{P}$ visible from y (resp., z). To achieve this, we choose y (resp., z) to be a point beyond the facet $\mathcal{P}_1$ (resp., $\mathcal{P}_m$) of $\mathcal{P}$, and beneath every other facet of $\mathcal{P}$. Let Q be the set of points consisting of y , z and the vertices of $\mathcal{P}$, and let $\mathcal{Q} = CH_{d+1}(Q)$ (refer to Fig. 2). Observe that the faces of $\mathcal{P}$, except for the facets $\mathcal{P}_1$ and $\mathcal{P}_m$, are all faces of $\mathcal{Q}$. To see that, notice that a supporting hyperplane H_F for a facet $F \in \mathcal{P}$, with $F \neq \mathcal{P}_1, \mathcal{P}_m$, is also a supporting hyperplane for $\mathcal{Q}$. Indeed, the vertices of F are vertices in $Q \setminus \{y, z\} \equiv P$, and thus, every vertex in P that is not a vertex of F strictly satisfies all hyperplane inequalities for $\mathcal{P}$. Also, by construction, the points y and z strictly satisfy all hyperplane inequalities apart from those for Π_1 and Π_m , respectively. Since H_F is a hyperplane other than Π_1 and Π_m we deduce that all points in $P \cup \{y, z\}$ lie on the same halfspace defined by H_F , and therefore H_F supports $\mathcal{Q}$. The faces of $\mathcal{Q}$ that are not faces of $\mathcal{P}$ are the faces in the star $\mathcal{S}_y$ of y and the star $\mathcal{S}_z$ of z . To verify this, consider a k -face F of $\mathcal{P}_1$, and let F_1 be a face in $\mathcal{P}_1$ that contains F . Let H_1 be a supporting hyperplane of F_1 with respect to $\mathcal{P}$. Tilt H_1 until it hits the point y , while keeping H_1 incident to F_1 , and call H'_1 this tilted hyperplane. H'_1 is a supporting hyperplane for y and the vertex set of $\mathcal{P}_1$, and thus is a supporting hyperplane for $\mathcal{Q}$. The same argument can be applied for the star of z . In fact, the boundary complex $\partial \mathcal{P}_1$ (resp., $\partial \mathcal{P}_m$) of $\mathcal{P}_1$ (resp., $\mathcal{P}_m$) is nothing but the link of y (resp., z) in $\mathcal{Q}$.




Figure 2: The polytope $\mathcal{Q}$. $\mathcal{Q}$ is constructed by adding a point y below Π_1 and a point z above Π_m , such the only vertices of $\mathcal{P}$ visible from y (resp., z) are the vertices of $\mathcal{P}_1$ (resp., $\mathcal{P}_m$).

It is easy to realize that, for $k < d$, the set of k -faces of $\mathcal{Q}$ is the union of the k -faces of $\mathcal{P}$, $\mathcal{S}_y$ and $\mathcal{S}_z$, where we have double-counted the k -faces of $\partial\mathcal{P}_1$ and $\partial\mathcal{P}_m$. For $k = d$ the set of facets of $\mathcal{Q}$ is the union of the set of facets of $\mathcal{S}_y$, $\mathcal{S}_z$ and $\mathcal{P}$, except for its two facets $\mathcal{P}_1$ and $\mathcal{P}_m$. We can concisely write the relations described above as:

$$f_k(\mathcal{Q}) = f_k(\mathcal{P}) + f_k(\mathcal{S}_y) + f_k(\mathcal{S}_z) - f_k(\mathcal{P}_1) - f_k(\mathcal{P}_m), \quad 0 \leq k \leq d. \quad (2)$$

where $f_d(\mathcal{P}_1) = f_d(\mathcal{P}_m) = 1$. The k -faces of $\mathcal{Q}$ in $\mathcal{S}_z$ are either k -faces of $\partial\mathcal{P}_m$ or k -faces defined by z and a $(k-1)$ -face of $\partial\mathcal{P}_m$. In fact, there exists a bijection between the $(k-1)$ -faces of $\partial\mathcal{P}_m$ and the k -faces of $\mathcal{S}_z$ containing z . Hence, we have:

$$f_k(\mathcal{S}_z) = f_k(\partial\mathcal{P}_m) + f_{k-1}(\partial\mathcal{P}_m), \quad 0 \leq k \leq d, \quad (3)$$

where $f_{-1}(\partial\mathcal{P}_m) = 1$ and $f_d(\partial\mathcal{P}_m) = 0$. Analogously, the k -faces of $\mathcal{Q}$ in $\mathcal{S}_y$ are either k -faces of $\partial\mathcal{P}_1$ or k -faces defined by y and a $(k-1)$ -face of $\partial\mathcal{P}_1$. As for $\mathcal{S}_z$, there exists a bijection between the $(k-1)$ -faces of $\partial\mathcal{P}_1$ and the k -faces of $\mathcal{S}_y$ containing y . Hence, we have:

$$f_k(\mathcal{S}_y) = f_k(\partial\mathcal{P}_1) + f_{k-1}(\partial\mathcal{P}_1), \quad 0 \leq k \leq d, \quad (4)$$

where $f_{-1}(\partial\mathcal{P}_1) = 1$ and $f_d(\partial\mathcal{P}_1) = 0$. Therefore, relation (2) can be rewritten as:

$$f_k(\mathcal{Q}) = f_k(\mathcal{P}) + f_k(\partial\mathcal{P}_1) + f_{k-1}(\partial\mathcal{P}_1) - f_k(\mathcal{P}_1) + f_k(\partial\mathcal{P}_m) + f_{k-1}(\partial\mathcal{P}_m) - f_k(\mathcal{P}_m), \quad 0 \leq k \leq d.$$

or, in the much simpler form:

$$f_k(\mathcal{Q}) = f_k(\mathcal{P}) + f_{k-1}(\partial\mathcal{P}_1) + f_{k-1}(\partial\mathcal{P}_m) - \alpha_k, \quad 0 \leq k \leq d, \quad (5)$$

where $\alpha_k = 0$, for $0 \leq k < d$, whereas $\alpha_d = 2$.

Let us now turn our attention on deriving bounds for the face numbers $f_k(\mathcal{Q})$. Since $\partial\mathcal{P}_\ell$, $\ell = 1, m$, is the boundary complex of the n_ℓ -vertex d -polytope $\mathcal{P}_\ell$, we have:

$$f_k(\partial\mathcal{P}_\ell) = O(n_\ell^{\min\{k+1, \lfloor \frac{d}{2} \rfloor\}}), \quad 0 \leq k \leq d, \quad \ell = 1, m. \quad (6)$$

Combining the bounds from Lemma 4 with relations (5) and (6), we get the following bounds for the lower-dimensional face numbers of $\mathcal{Q}$:

$$f_k(\mathcal{Q}) = O\left(\sum_{1 \leq i \neq j \leq m} n_i n_j^k + \sum_{i=1}^m n_i^{\min\{k+1, \lfloor \frac{d}{2} \rfloor\}}\right) = O\left(\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor}\right), \quad 0 \leq k \leq \lfloor \frac{d}{2} \rfloor. \quad (7)$$

However, $\mathcal{Q}$ is a simplicial $(d+1)$ -polytope, since the facets of $\mathcal{P}$ are simplicial, and since all facets of $\mathcal{S}_y$ and $\mathcal{S}_z$ are simplicial (the facets of $\mathcal{S}_y$ and $\mathcal{S}_z$ are defined via simplicial $(d-1)$ -faces of $\mathcal{P}$ and the points y and z , respectively). Let us recall the defining equations for the elements of the h -vector of $\mathcal{Q}$ in terms of the elements of the f -vector of $\mathcal{Q}$:

$$h_k(\mathcal{Q}) = \sum_{i=0}^k (-1)^{k-i} \binom{d+1-i}{d+1-k} f_{i-1}(\mathcal{Q}), \quad 0 \leq k \leq d+1. \quad (8)$$

Combining equations (8) with relations (7), as well as the fact that $f_{-1}(\mathcal{Q}) = 1$, we get:

$$h_k(\mathcal{Q}) = O\left(\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor}\right), \quad 0 \leq k \leq \lfloor \frac{d+1}{2} \rfloor. \quad (9)$$

We are now going to use the Dehn-Sommerville equations for $\mathcal{Q}$ to bound the number of k -faces $f_k(\mathcal{Q})$ of $\mathcal{Q}$, for $k \geq \lfloor \frac{d+1}{2} \rfloor$. The Dehn-Sommerville equations can be rewritten as follows (cf. [28, Section 8.4]):

$$f_{k-1}(\mathcal{Q}) = \sum_{i=0}^{\frac{d+1}{2}} \left(\binom{d+1-i}{k-i} + \binom{i}{k-d-1+i} \right) h_i(\mathcal{Q}), \quad \lfloor \frac{d+1}{2} \rfloor \leq k \leq d+1, \quad (10)$$

where the symbol $\sum_{i=0}^{\frac{\delta}{2}*}$ denotes the sum where the last term is halved if and only if the quantity $\frac{\delta}{2}$ is integral (which is our case since in our setting $\delta = d + 1$, which is even). Combining relations (9) and (10), we get:

$$f_{k-1}(\mathcal{Q}) = O\left(\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor}\right), \quad \lfloor \frac{d+1}{2} \rfloor \leq k \leq d+1. \quad (11)$$

Now using relation (5), we arrive at the following bounds on the number of k -faces of $\mathcal{P}$ for $\lfloor \frac{d+1}{2} \rfloor \leq k \leq d$:

$$f_k(\mathcal{P}) = f_k(\mathcal{Q}) - f_{k-1}(\partial\mathcal{P}_1) - f_{k-1}(\partial\mathcal{P}_m) + \alpha_k \leq f_k(\mathcal{Q}) = O\left(\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor}\right),$$

where we used the fact that $f_{d-1}(\partial\mathcal{P}_1) > \alpha_d$, since $f_{d-1}(\partial\mathcal{P}_1) \geq d+1 \geq 4$. $\square$

By Lemmas 4 and 5, we deduce that the worst-case complexity of the convex hull $CH_{d+1}(\mathcal{P})$ is $O(\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor})$. As we will see in Subsection 4.2 (see Corollary 11), this bound is asymptotically tight for any odd $d \geq 3$. Hence:

Theorem 6. *Let $\mathcal{P} = \{\mathcal{P}_1, \mathcal{P}_2, \dots, \mathcal{P}_m\}$ be a set of a fixed number of $m \geq 2$ d -polytopes, lying on m parallel hyperplanes of $\mathbb{E}^{d+1}$, where $d \geq 3$ and d is odd. The worst-case complexity of the convex hull $CH_{d+1}(\mathcal{P})$ is $\Theta(\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor})$, where $n_i = f_0(\mathcal{P}_i)$, $1 \leq i \leq m$.*

3 Computing the convex hull of parallel convex polytopes

The upper bound on the worst-case complexity of $CH_{d+1}(\mathcal{P})$ in Theorem 6 suggests that, in order to compute $CH_{d+1}(\mathcal{P})$ for $d \geq 3$ odd, it pays off to use an output-sensitive algorithm for constructing the convex hull of the point set P formed by the vertices of the $\mathcal{P}_i$'s.

Let us briefly discuss the output-sensitive algorithms applicable in our setting; in what follows n_i is the number of vertices of $\mathcal{P}_i$, n is the total number of input points and f the size of the output. The dimension $d+1 \geq 4$ below is the dimension of the ambient space $\mathbb{E}^{d+1}$ and is considered to be even; the polytopes $\mathcal{P}_i$ are then d -dimensional. One of the earliest algorithms is Seidel's shelling algorithm [27] that runs in $O(n^2 + f \log n)$ time. The preprocessing step of Seidel's algorithm was later on improved by Matoušek and Schwarzkopf [21], resulting in an $O(n^{2-2/(\lfloor (d+1)/2 \rfloor + 1)} + f \log n)$ time algorithm, for any fixed $\epsilon > 0$. Chan, Snoeyink and Yap [11] describe a divide-and-conquer algorithm for constructing four-dimensional convex hulls in $O((n+f) \log^2 f)$ time. Finally, Chan [10] improved the gift-wrapping algorithm of Chand and Kapur [12], yielding an $O(n \log f + (nf)^{1-1/(\lfloor (d+1)/2 \rfloor + 1)} \log^{O(1)} n)$ time algorithm.

In our case we have $f = O(\sum_{1 \leq i \neq j \leq m} n_i n_j^{\alpha-1})$, where $\alpha = \lfloor \frac{d+1}{2} \rfloor \geq 2$ (recall that we consider only odd d). Applying the algorithms above we get the following. Seidel's algorithm runs in $O(n^2 + (\sum_{1 \leq i \neq j \leq m} n_i n_j^{\alpha-1}) \log n)$ time. Matoušek and Schwarzkopf's improvement results in an $O(n^{2-2/(\alpha+1)+\epsilon} + (\sum_{1 \leq i \neq j \leq m} n_i n_j^{\alpha-1}) \log n)$ time algorithm. Chan's algorithm yields a running time of $O((\sum_{1 \leq i \neq j \leq m} n_i^{\alpha/(\alpha+1)} n_j^{\alpha^2/(\alpha+1)}) \log^{O(1)} n)$. Finally, the algorithm by Chan, Snoeyink and Yap gives $O((\sum_{1 \leq i \neq j \leq m} n_i n_j) \log^2 n)$. For all $d \geq 3$, i.e., for $\alpha \geq 2$, Chan's algorithm is never faster than the other two algorithms. For $d \geq 5$, i.e., $\alpha \geq 3$, it makes no difference to choose between the algorithm of Seidel and that of Matoušek and Schwarzkopf. For $d = 3$ the situation is a bit more complicated. For some choices for the n_i 's Matoušek and Schwarzkopf's algorithm outperforms all

other algorithms, whereas for other choices of the n_i 's the algorithm of Chan, Snoeyink and Yap is the best choice. Consider, for example, the cases:

- (1) $n_1 = \Theta(n)$, $n_2 = o(\frac{n^{1/3+\epsilon}}{\log^2 n})$ and $n_i = O(1)$ for $3 \leq i \leq m$, and
- (2) $n_1 = \Theta(n)$, $n_2 = \omega(\frac{n^{1/3+\epsilon}}{\log n})$ and $n_i = O(1)$ for $3 \leq i \leq m$.

In the first case Chan, Snoeyink and Yap's algorithm runs in $o(n^{4/3+\epsilon})$ time and outperforms all other algorithms. In the second case, Matoušek and Schwarzkopf's algorithm seems to be the best choice, since its running time is $O(n_2 n \log n)$ (compared to Chan, Snoeyink and Yap's algorithm, the running time of which is $O(n_2 n \log^2 n)$). Summarizing, we arrive at the following:

Theorem 7. *Let $\mathcal{P} = \{\mathcal{P}_1, \mathcal{P}_2, \dots, \mathcal{P}_m\}$ be a set of m d -polytopes, lying on m parallel hyperplanes of $\mathbb{E}^{d+1}$, where $d \geq 3$ and d odd. Let $n_i = f_0(\mathcal{P}_i)$, $1 \leq i \leq m$, and $n = \sum_{i=1}^m n_i$. We can compute the convex hull $CH_{d+1}(\mathcal{P})$ in $O((\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor}) \log n)$ time for $d \geq 5$, and $O(\min\{n^{4/3+\epsilon} + (\sum_{1 \leq i \neq j \leq m} n_i n_j) \log n, (\sum_{1 \leq i \neq j \leq m} n_i n_j) \log^2 n\})$ time for $d = 3$, for any fixed $\epsilon > 0$.*

4 Convex hulls of spheres with a fixed number of distinct radii

In this section we derive tight upper and lower bounds on the worst-case complexity of the convex hull of a set of spheres in $\mathbb{E}^d$ having m distinct radii, where m is considered to be fixed.

Let Σ be a set of n spheres $\sigma_k = (c_k, r_k)$, $1 \leq k \leq n$, in $\mathbb{E}^d$, and let $CH_d(\Sigma)$ be the convex hull of the spheres in Σ . A *face of circularity* ℓ of $CH_d(\Sigma)$, $0 \leq \ell \leq d-1$, is a maximal connected portion of the boundary of $CH_d(\Sigma)$ consisting of points where the supporting hyperplanes are tangent to a given set of $(d-\ell)$ spheres of Σ . In the special case where all spheres have the same radius, $CH_d(\Sigma)$ is combinatorially equivalent to the convex hull $CH_d(K)$ of the centers K of spheres in Σ , in the sense that each face of circularity ℓ of $CH_d(\Sigma)$ corresponds to a unique $(d-\ell-1)$ -face of $CH_d(K)$, for $0 \leq \ell \leq d-1$.

We consider here the case where the radii r_k can take m distinct values, i.e., $r_k \in \{\rho_1, \rho_2, \dots, \rho_m\}$. Without loss of generality we may assume that $0 < \rho_1 < \rho_2 < \dots < \rho_m$. We identify $\mathbb{E}^d$ with the hyperplane $H_0 = \{x_{d+1} = 0\}$ of $\mathbb{E}^{d+1}$ and we call the $(d+1)$ -axis of $\mathbb{E}^{d+1}$ the *vertical axis*, while the expression *above* will refer to the $(d+1)$ -coordinate. Let Π_i , $1 \leq i \leq m$, be the hyperplane $\{x_{d+1} = \rho_i\}$ in $\mathbb{E}^{d+1}$, and let P be the point set in $\mathbb{E}^{d+1}$ obtained by mapping each sphere σ_k to the point $p_k = (c_k, r_k)$. Let P_i denote the subset of P containing points that belong to the hyperplane Π_i , and let n_i be the cardinality of P_i . We denote by $\mathcal{P}$ the convex hull of the points in P (i.e., $\mathcal{P} = CH_{d+1}(P)$). We further denote by $\mathcal{P}_i$ the convex hull of the points in P_i (i.e., $\mathcal{P}_i = CH_d(P_i)$); more precisely, we identify Π_i with $\mathbb{E}^d$, and then define $\mathcal{P}_i$ to be the convex hull of the points in P_i , seen as points in $\mathbb{E}^d$. We use $\mathcal{P}$ to denote the set of the $\mathcal{P}_i$'s. Let $\hat{P}_i$ be the subset of P_i that defines $\mathcal{P}_i$ (i.e., the points in $\hat{P}_i$ are the vertices of $\mathcal{P}_i$ and thus $\mathcal{P}_i = CH_d(\hat{P}_i)$), and let $\hat{n}_i \leq n_i$ be the cardinality of $\hat{P}_i$. Finally, let $\hat{P} = \bigcup_{i=1}^m \hat{P}_i$ and $\hat{\mathcal{P}} = CH_{d+1}(\hat{P})$. Notice that it is possible that $\mathcal{P} \neq \hat{\mathcal{P}}$; such a situation will arise if $P_1 \neq \hat{P}_1$ (resp., $P_m \neq \hat{P}_m$), in which case the intersection of $\mathcal{P}$ with Π_1 (resp., Π_m) will consist of more than one d -face of $\mathcal{P}$. On the other hand $\mathcal{P}$ and $\hat{\mathcal{P}}$ have the same interior.

Let λ_0 be the half lower hypercone in $\mathbb{E}^{d+1}$ with arbitrary apex, vertical axis, and angle at the apex equal to $\frac{\pi}{4}$. By $\lambda(p)$ we denote the translated copy of λ_0 with apex at p ; observe that the intersection of the hypercone $\lambda(p_k)$ with the hyperplane H_0 is identical to the sphere σ_k . Let Λ be the set of the lower half hypercones $\{\lambda(p_1), \lambda(p_2), \dots, \lambda(p_n)\}$ in $\mathbb{E}^{d+1}$ associated with the spheres of Σ . The intersection of the convex hull $CH_{d+1}(\Lambda)$ with H_0 is equal to $CH_d(\Sigma)$.

Let O' be a point in the interior of $\mathcal{P}$. We then have the following:

Theorem 8 ([6, Theorem 1]). *Any hyperplane of $\mathbb{E}^d$ supporting $CH_d(\Sigma)$ is the intersection with H_0 of a unique hyperplane H of $\mathbb{E}^{d+1}$ satisfying the following three properties:*

1. *H supports $\mathcal{P}$,*
2. *H is the translated copy of a hyperplane tangent to λ_0 along one of its generatrices,*
3. *H is above O' .*

Conversely, let H be a hyperplane of $\mathbb{E}^{d+1}$ satisfying the above three properties. Its intersection with H_0 is a hyperplane of $\mathbb{E}^d$ supporting $CH_d(\Sigma)$.

Theorem 8 implies an injection $\varphi : CH_d(\Sigma) \rightarrow \mathcal{P}$ that maps each face of circularity $(d - \ell - 1)$ of $CH_d(\Sigma)$ to a unique ℓ -face of $\mathcal{P}$, for $0 \leq \ell \leq d - 1$. Theorem 8 also implies that points in $P_i \setminus \hat{P}_i$, $1 \leq i \leq m$, can never be points on a supporting hyperplane H of $\mathcal{P}$ satisfying the three properties of the theorem. Therefore, φ is, in fact, an injection that maps each face of circularity $(d - \ell - 1)$ of $CH_d(\Sigma)$ to a unique ℓ -face of $\hat{\mathcal{P}}$, $0 \leq \ell \leq d - 1$. Observe that $\hat{\mathcal{P}}$ is the convex hull of the set $\mathcal{P}$ of m convex polytopes lying on m parallel hyperplanes of $\mathbb{E}^{d+1}$. By employing Theorem 6 of Section 2, we deduce that $\hat{\mathcal{P}}$'s complexity is $O(\sum_{1 \leq i \neq j \leq m} \hat{n}_i \hat{n}_j^{\lfloor \frac{d}{2} \rfloor}) = O(\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor})$, which, via the injection $\varphi : CH_d(\Sigma) \rightarrow \hat{\mathcal{P}}$, is also an upper bound for the worst-case complexity of $CH_d(\Sigma)$.

4.1 Lower bound construction with two distinct radii

For any even dimension 2δ , the trigonometric moment curve $\gamma_{2\delta}^{tr}(t)$ in $\mathbb{E}^{2\delta}$ is the curve:

$$\gamma_{2\delta}^{tr}(t) = (\cos t, \sin t, \cos 2t, \sin 2t, \dots, \cos \delta t, \sin \delta t), \quad t \in [0, \pi).$$

Notice that points on $\gamma_{2\delta}^{tr}(t)$ are points on the sphere of $\mathbb{E}^{2\delta}$ centered at the origin with radius equal to $\sqrt{\delta}$. For any set P of n points on $\gamma_{2\delta}^{tr}(t)$, the convex hull $CH_{2\delta}(P)$ is a polytope $\mathcal{Q}$ combinatorially equivalent to the cyclic polytope $C_{2\delta}(n)$ (cf. [17, 28]). Therefore, $f_{2\delta-1}(\mathcal{Q}) = \Theta(n^\delta)$.

Suppose now that the ambient space is $\mathbb{E}^d$, where $d \geq 3$ is odd. Let H_1 be H_2 be the hyperplanes $\{x_d = z_1\}$ and $\{x_d = z_2\}$, where $z_1, z_2 \in \mathbb{R}$ and $z_2 > z_1 + 2(n_2 + 2)\sqrt{\delta}$; the quantity n_2 will be defined below. Consider a set Σ_1 of $n_1 + 1$ points, treated as spheres of $\mathbb{E}^d$ of zero radius, on the $(d - 1)$ -dimensional trigonometric moment curve $\gamma_{d-1}^{tr}(t)$ embedded in H_1 (please refer to Fig. 3(left), as well as Fig. 4 for the view of the construction from the positive x_d -axis). Among the $n_1 + 1$ points, the first n_1 points are chosen with $t \in (0, \frac{\pi}{2})$, whereas for the remaining point we require that $t \in (\frac{\pi}{2}, \pi)$. This implies that the x_1 -coordinate of the first n_1 points of Σ_1 is positive, whereas the x_1 -coordinate of the last point of Σ_1 is negative. Let Σ_2 be the projection, along the x_d -axis, of Σ_1 on the hyperplane H_2 . Clearly, the $n_1 + 1$ points of Σ_2 in H_2 lie on the $(d - 1)$ -dimensional trigonometric moment curve $\gamma_{d-1}^{tr}(t)$ embedded in H_2 . The points of Σ_i , $i = 1, 2$, lie on a $(d - 2)$ -dimensional sphere of $\mathbb{E}^d$, centered at the point $(0, 0, \dots, 0, z_i)$, with radius $\sqrt{\delta}$. Moreover, the number of facets of the polytope $\mathcal{Q}_i = CH_{d-1}(\Sigma_i)$ is $\Theta(n_1^{\lfloor \frac{d-1}{2} \rfloor}) = \Theta(n_1^{\lfloor \frac{d}{2} \rfloor})$. The convex hull of the $2(n_1 + 1)$ points of $\Sigma_1 \cup \Sigma_2$ is a prism Δ . Δ consists of $\Theta(n_1^{\lfloor \frac{d}{2} \rfloor})$ facets not lying on H_1 or H_2 , called the *vertical facets* of the prism, the $(d - 1)$ -face of $CH_{d-1}(\Sigma_1)$, called the *bottom facet*, and the $(d - 1)$ -face of $CH_{d-1}(\Sigma_2)$, called the *top facet*. For each vertical facet F of Δ , we denote by $\vec{v}_F$ the unit normal vector of F pointing outside Δ , and by F^+ (resp., F^-) the positive (resp., negative) open halfspace delimited by the supporting hyperplane of F . Regarding the ridges of Δ , those that are intersections of vertical facets of Δ will be referred to as *vertical ridges*. Notice that the vertical ridges of Δ are perpendicular to H_1 and H_2 .

Let Y be the oriented hyperplane $\{x_1 = 0\}$ with unit normal vector $\vec{v} = (1, 0, \dots, 0)$. Let also Y^+ and Y^- be the positive and negative open halfspaces of $\mathbb{E}^d$ delimited by Y , respectively. Y contains the x_d -axis, and is perpendicular to the hyperplanes H_1 and H_2 . Recall that n_1 points

of Σ_i are contained in Y^+ , whereas exactly one point of Σ_i is contained in Y^- . Clearly, Y is in general position with respect to Δ , $\mathcal{Q}_1$ and $\mathcal{Q}_2$. Let $\tilde{\mathcal{Q}}_i$ be the intersection of $\mathcal{Q}_i$ with Y , and let $\mathcal{F}_i$ be the set of faces of $\mathcal{Q}_i$ intersected by Y . $\tilde{\mathcal{Q}}_i$ is a $(d-2)$ -polytope, and its number of vertices is at most n_1 , since Y cuts at most n_1 edges of $\mathcal{Q}_i$. This implies that the complexity of $\tilde{\mathcal{Q}}_i$ is $O(n_1^{\lfloor \frac{d-2}{2} \rfloor}) = O(n_1^{\lfloor \frac{d}{2} \rfloor - 1})$; the same bound holds for $\mathcal{F}_i$. Since there are no facets of Δ in Y^- (Y^- contains a single vertex of $\mathcal{Q}_i$), and since the number of facets of $\mathcal{Q}_i$ is $\Theta(n_1^{\lfloor \frac{d}{2} \rfloor})$, we conclude that the number of facets of $\mathcal{Q}_i$ contained in Y^+ is also $\Theta(n_1^{\lfloor \frac{d}{2} \rfloor})$; the same bound holds for the number of vertical facets of Δ in Y^+ .

Define now a set $\Sigma_3 = \{\sigma_0, \sigma_1, \dots, \sigma_{n_2+1}\}$ of $n_2 + 2$ spheres in $\mathbb{E}^d$, where $\sigma_k = (c_k, \rho)$, and $c_k = (0, \dots, 0, (2k+1)\sqrt{\delta})$, $0 \leq k \leq n_2 + 1$. In other words, the sphere σ_k is centered on the x_d -axis, with the d -th coordinate of its center c_k being $(2k+1)\sqrt{\delta}$, while its radius is ρ . We choose ρ to be smaller than $\sqrt{\delta}$, but large enough so that each sphere σ_i satisfies the following two properties:

- (1) it does not intersect any of the ridges of Δ (including the vertical ridges of Δ), and
- (2) it intersects the interior of all vertical facets of Δ .

Notice also that for this choice for ρ , none of the spheres in Σ_3 intersects the hyperplanes H_1 and H_2 (recall that $z_2 > z_1 + 2(n_2 + 2)\sqrt{\delta}$), while the spheres in Σ_3 are pairwise disjoint; these two observations, however, are not critical for our construction.

We are now going to perturb the centers of the spheres in Σ_3 to get a new set of spheres Σ'_3 (see Fig. 3(right), as well as Fig. 5 for the view of the construction from the positive x_d -axis). Define σ'_k to be the sphere with radius ρ and center $c'_k = c_k + (\sum_{\ell=0}^k \frac{\varepsilon}{2^\ell})\vec{v} = c_k + \varepsilon(2 - \frac{1}{2^k})\vec{v}$, where $0 < \varepsilon \ll 1$. The quantity ε is chosen so that the spheres in Σ'_3 satisfy almost the same conditions as the spheres in Σ_3 . In particular, we require that condition (1) is still satisfied, while we relax the requirement on condition (2): we now require that σ'_k intersects the interior of all vertical facets of Δ contained in Y^+ . In addition to the two conditions above, we also require that for each k , $0 \leq k \leq n_2 + 1$, the $(d-2)$ -dimensional sphere $\sigma_k \cap \sigma'_k$ is contained in F^- for all vertical facets F of Δ in Y^+ .

We will now show that for each pair (σ'_k, F) , where $1 \leq k \leq n_2$ and F is a vertical facet of Δ in Y^+ , the spherical cap $F^+ \cap \sigma'_k$ induces a face of circularity $(d-1)$ in $CH_d(\Sigma)$. Let F_1 and F_2 be the ridges of Δ on the boundary of F contained in the top and bottom facet, respectively. Finally, let S_k be the supporting hyperplane of σ_k parallel to F ; we consider S_k to be oriented as F (i.e., the unit normal vector of S_k is $\vec{v}_F$), and thus σ_k lies in the closure of the negative halfspace delimited by S_k . Notice that S_k is also a supporting hyperplane for Σ_3 . Let S'_k be the hyperplane we get by translating S_k by the vector $\varepsilon(2 - \frac{1}{2^k})\vec{v}$. S'_k supports σ'_k , but fails to be a supporting hyperplane for Σ'_3 . More precisely, S'_k intersects all spheres σ'_j with $j > k$, whereas all spheres σ'_j with $j < k$, are contained in the negative open halfspace delimited by S'_k . We can, however, perturb S'_k so that it supports Σ'_3 : simply slide S'_k on sphere σ'_k towards F_1 , while maintaining the property that it remains parallel to F_1 and F_2 . We keep sliding S'_k until it has empty intersection with any sphere σ'_j with $j > k$. Notice that due to the way we have perturbed the centers of the spheres in Σ_3 to get Σ'_3 , the new hyperplane S''_k we get via this transformation is a supporting hyperplane for Σ'_3 . In fact, S''_k is a supporting hyperplane for the sphere set $\Sigma = \Sigma_1 \cup \Sigma_2 \cup \Sigma'_3$ (it touches $CH_d(\Sigma)$ at σ'_k only), which implies that S''_k corresponds to a unique face of circularity $(d-1)$ on $CH_d(\Sigma)$.

The same construction can be done for all k with $1 \leq k \leq n_2$, and for all vertical facets of Δ in Y^+ . Since we have $\Theta(n_1^{\lfloor \frac{d}{2} \rfloor})$ vertical facets of Δ in Y^+ , we can construct $n_2\Theta(n_1^{\lfloor \frac{d}{2} \rfloor})$ distinct supporting hyperplanes of $CH_d(\Sigma)$, corresponding to distinct faces of circularity $(d-1)$ on $CH_d(\Sigma)$. Hence the complexity of $CH_d(\Sigma)$ is $\Omega(n_2n_1^{\lfloor \frac{d}{2} \rfloor})$. Without loss of generality, we may assume that $n_2 \leq n_1$, in which case we have $n_2n_1^{\lfloor \frac{d}{2} \rfloor} \geq \frac{1}{2}(n_2n_1^{\lfloor \frac{d}{2} \rfloor} + n_1n_2^{\lfloor \frac{d}{2} \rfloor})$. Hence, we arrive at the following:




Figure 3: The lower bound construction in the case of two radii. The points in Σ_1 (resp., Σ_2) are shown in black (resp., white). The hyperplane Y is shown in green, while the prism Δ is shown in black. The facet F in blue is one of the vertical facets of Δ in Y^+ . The sphere sets Σ_3 (left) and Σ'_3 (right) are shown in red. The red spherical caps on the left correspond to a unique supporting hyperplane of $CH_d(\Sigma_1 \cup \Sigma_2 \cup \Sigma_3)$. The red spherical caps on the right correspond to faces of $CH_d(\Sigma_1 \cup \Sigma_2 \cup \Sigma'_3)$ of circularity $(d-1)$.




Figure 4: View from the positive x_d -axis of the construction in Fig. 3(left). The silhouettes of all spheres in Σ_3 coincide.




Figure 5: View from the positive x_d -axis of the construction in Fig. 3(right).

Theorem 9. Fix some odd $d \geq 3$. There exists a set Σ of spheres in $\mathbb{E}^d$, consisting of n_i spheres of radius ρ_i , $i = 1, 2$, with $\rho_1 \neq \rho_2$, such that the complexity of the convex hull $CH_d(\Sigma)$ is $\Omega(n_1 n_2^{\lfloor \frac{d}{2} \rfloor} + n_2 n_1^{\lfloor \frac{d}{2} \rfloor})$.

4.2 Lower bound construction with m distinct radii

We can easily generalize the lower bound construction of the previous subsection in the case where we have n_i spheres of radius ρ_i , $1 \leq i \leq m$, $m \geq 3$, and the radii ρ_i are considered to be mutually distinct.

As in the previous subsection, the ambient space is $\mathbb{E}^d$, where $d \geq 3$ is odd. Let $N_1 = \sum_{i=2}^m n_i$ and $N_2 = n_1$. We construct the set $\Sigma = \Sigma_1 \cup \Sigma_2 \cup \Sigma'_3$ as in the previous subsection where Σ_1 and Σ_2 contain each $N_1 + 1$ points and Σ'_3 contains $N_2 + 2$ spheres of some appropriate radius ρ (recall that in the construction of the previous subsection $\rho \approx \sqrt{\frac{d-1}{2}} \geq 1$). We then replace n_i among the N_1 points of Σ_1 (resp., Σ_2) contained in Y^+ by spheres with the same center and radius equal to r^i , where $0 < r \ll 1$. Furthermore, we replace the unique point of Σ_1 (resp., Σ_2) in Y^- by a sphere of the same center and radius r^2 . We choose r small enough so that the following two conditions hold:

- (1) the prism $\Delta_r = CH_d(\Sigma_1 \cup \Sigma_2)$ is combinatorially equivalent² to the prism Δ_0 (this is the prism we get for $r = 0$, which is the prism Δ of the previous subsection), and
- (2) the two requirements for the spheres in Σ'_3 are still satisfied: each σ'_k does not intersect any of the ridges of Δ_r , while each σ'_k intersects the interior of all vertical facets of Δ_r contained in Y^+ .

As described in the previous subsection, the convex hull $CH_d(\Sigma)$ of the set $\Sigma = \Sigma_1 \cup \Sigma_2 \cup \Sigma'_3$ of $2(N_1 + 1) + N_2 + 2$ spheres has $N_2 \Theta(N_1^{\lfloor \frac{d}{2} \rfloor})$ faces of circularity $(d - 1)$, and hence its complexity is $\Omega(N_2 N_1^{\lfloor \frac{d}{2} \rfloor}) = \Omega(n_1 (\sum_{i=2}^m n_i)^{\lfloor \frac{d}{2} \rfloor})$. Without loss of generality we may assume that $n_2 \geq n_1 \geq n_i$ for all $3 \leq i \leq m$, in which case we have: $n_1 (\sum_{i=2}^m n_i)^{\lfloor \frac{d}{2} \rfloor} \geq n_1 n_2^{\lfloor \frac{d}{2} \rfloor} \geq \frac{1}{m(m-1)} (\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor})$. Since m is fixed, we conclude that the complexity of $CH_d(\Sigma)$ is $\Omega(\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor})$.

Theorem 10. Fix some odd $d \geq 3$. There exists a set Σ of spheres in $\mathbb{E}^d$, consisting of n_i spheres of radius ρ_i , $1 \leq i \leq m$, with $\rho_1 < \rho_2 < \dots < \rho_m$ and $m \geq 3$ fixed, such that the complexity of the convex hull $CH_d(\Sigma)$ is $\Omega(\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor})$.

Consider again the injection $\varphi : CH_d(\Sigma) \rightarrow \hat{\mathcal{P}}$. We have shown above that the worst-case complexity of $CH_d(\Sigma)$ is $\Omega(\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor})$, when $d \geq 3$ is odd and $m \geq 2$ is fixed. Since φ is injective, this lower bound also applies to the complexity of $\hat{\mathcal{P}}$. This establishes our lower bound claim in Theorem 6:

Corollary 11. Let $\mathcal{P} = \{\mathcal{P}_1, \mathcal{P}_2, \dots, \mathcal{P}_m\}$ be a set of m d -polytopes, lying on m parallel hyperplanes of $\mathbb{E}^{d+1}$, with $d \geq 3$, d odd, and both d and m are fixed. The worst-case complexity of $CH_{d+1}(\mathcal{P})$ is $\Omega(\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor})$, where $n_i = f_0(\mathcal{P}_i)$, $1 \leq i \leq m$.

²Combinatorial equivalence here means that each face of circularity ℓ of Δ_r corresponds to a unique $(d - \ell - 1)$ -face of Δ_0 .

5 Computing convex hulls of spheres

In this section we focus our attention to the computation of the convex hull $CH_d(\Sigma)$ of Σ . We use the same notation as in Section 4.

Polarity with respect to the point O' is a one-to-one transformation which maps points of $\mathbb{E}^{d+1}$ distinct from O' to hyperplanes of $\mathbb{E}^{d+1}$ that do not contain O' (e.g., see [8, Section 7.1.3]). For a point q in $\mathbb{E}^{d+1}$ distinct from O' , we define its *polar hyperplane* $q^\star$ by

$$q^\star = \{X \in \mathbb{E}^{d+1} \mid q \cdot X = 1\},$$

while for a hyperplane H of $\mathbb{E}^{d+1}$ not containing O' , the *polar point* or *pole* $H^\star$ of H is defined by

$$H^\star \cdot X = 1, \quad \forall X \in H.$$

The *polar set* of a set of hyperplanes is the set of the poles of these hyperplanes. Finally, we denote by H^- the closed halfspace bounded by H and containing O' .

Consider again the point set P in $\mathbb{E}^{d+1}$ that we get by lifting the spheres of Σ . The polytope $\mathcal{P}^\star$ in $\mathbb{E}^{d+1}$ defined as the intersection of the halfspaces $p_k^{\star-}$, $1 \leq k \leq n$, is the *polar polytope* or *dual polytope* of $\mathcal{P}$. $\mathcal{P}$ and $\mathcal{P}^\star$ are dual in the sense that there is a bijection between the ℓ -faces of $\mathcal{P}$ and the $(d - \ell)$ -faces of $\mathcal{P}^\star$, which is inclusion-reversing: a hyperplane supporting $\mathcal{P}$ along an ℓ -face F , has its pole on the corresponding $(d - \ell)$ -face $F^\star$ of $\mathcal{P}^\star$. In a similar manner, we can define the polytope $\hat{\mathcal{P}}^\star$, which is the polar polytope of $\hat{\mathcal{P}}$.

Let $H_{O'}$ be the hyperplane parallel to H_0 passing through O' and let $H_{O'}^+$ denote the open halfspace of $\mathbb{E}^{d+1}$ above $H_{O'}$. Boissonnat *et al.* [6, Proposition] have shown that:

1. The polar set of the hyperplanes, which are translated copies of the hyperplanes tangent to λ_0 , is the hypercone K with apex at O' , vertical axis, and angle at the apex equal to $\frac{\pi}{4}$.
2. The polar set of the hyperplanes above O' is the halfspace $H_{O'}^+$.

An immediate consequence of these properties is that the polar set of the hyperplanes supporting $\mathcal{P}$, tangent to at least one hypercone of Λ along a generatrix, and above O' , is $\mathcal{P}^\star \cap K \cap H_{O'}^+$.

Boissonnat *et al.* [6] have used this property in order to propose an algorithm for computing $CH_d(\Sigma)$ in $O(n^{\lceil \frac{d}{2} \rceil} + n \log n)$ time for any $d \geq 2$. Below, we describe a slightly modified algorithm that takes into account the fact that the radii of the spheres in Σ can take a fixed number of $m \geq 2$ distinct values. Our algorithm consists of the following five steps:

1. For all i with $1 \leq i \leq m$: determine the set $P_i = P \cap \Pi_i$, construct the convex hull $\mathcal{P}_i = CH_d(P_i)$, and compute the set $\hat{P}_i$.
2. Compute the polytope $\hat{\mathcal{P}} = CH_{d+1}(\hat{P})$, and choose a point O' inside $\hat{\mathcal{P}}$.
3. Compute the polar polytope $\hat{\mathcal{P}}^\star$ of $\hat{\mathcal{P}}$ with respect to O' .
4. Compute the intersection between $\hat{\mathcal{P}}^\star$, the hypercone K , and the halfspace $H_{O'}^+$.
5. Compute the incidence graph of the facets in $CH_d(\Sigma)$ from the incidence graph of the faces of $\hat{\mathcal{P}}^\star$ intersecting K and $H_{O'}^+$.

Determining all the sets P_i takes $\Theta(n)$ time, whereas constructing the polytope $\mathcal{P}_i$ takes $O(n_i^{\lceil \frac{d}{2} \rceil} + n_i \log n_i)$ time. Determining $\hat{P}_i$ from the representation of $\mathcal{P}_i$ can easily be done in $O(n_i \log n_i)$ time. We thus conclude that step 1 of the algorithm takes $O(n^{\lceil \frac{d}{2} \rceil} + n \log n)$ time. Computing $\hat{\mathcal{P}}^\star$ from $\hat{\mathcal{P}}$ takes linear time in the size of $\hat{\mathcal{P}}$. Moreover, the complexity of computing $\hat{\mathcal{P}}^\star \cap K \cap H_{O'}^+$ is linear in the complexity of $\hat{\mathcal{P}}^\star$, and thus $\hat{\mathcal{P}}$, since K and $H_{O'}^+$ have constant complexity. The complexity for computing the incidence graph of $CH_d(\Sigma)$ from the incidence graph of $\hat{\mathcal{P}}^\star \cap K \cap H_{O'}^+$ is again linear in $\hat{\mathcal{P}}^\star$ and $\hat{\mathcal{P}}$. Therefore, the total time needed for steps 2–5 of the algorithm is a linear function of the time needed to compute $\hat{\mathcal{P}}$. Summarizing:

Theorem 12. Let Σ be a set of n spheres in $\mathbb{E}^d$, having m distinct radii $\rho_1, \rho_2, \dots, \rho_m$, with $d \geq 3$, d odd, and both d and m are fixed. Let n_i be the number of spheres in Σ with radius ρ_i , $1 \leq i \leq m$. We can compute the convex hull $CH_d(\Sigma)$ in $O(n^{\lfloor \frac{d}{2} \rfloor} + n \log n + T_{d+1}(n_1, n_2, \dots, n_m))$ time, where $T_{d+1}(n_1, n_2, \dots, n_m)$ stands for the time to compute the convex hull of m d -polytopes $\{\mathcal{P}_1, \mathcal{P}_2, \dots, \mathcal{P}_m\}$ lying on m parallel hyperplanes in $\mathbb{E}^{d+1}$, with $n_i = f_0(\mathcal{P}_i)$, $1 \leq i \leq m$.

As described in Section 3,

$$T_{d+1}(n_1, \dots, n_m) = O((\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor}) \log n),$$

for any odd $d \geq 5$, whereas for $d = 3$ we have

$$T_4(n_1, \dots, n_m) = O(\min\{n^{4/3+\epsilon} + (\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor}) \log n, (\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor}) \log^2 n\}).$$

Hence, we can compute the convex hull $CH_d(\Sigma)$ in $O((\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor}) \log n)$ time for any odd $d \geq 5$, and in $O(\min\{n^{4/3+\epsilon} + (\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor}) \log n, (\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor}) \log^2 n\})$ time for $d = 3$.

6 Summary and open problems

In this paper we have considered the problem of computing the worst-case complexity of the convex hull $CH_{d+1}(\mathcal{P})$ of a set $\mathcal{P} = \{\mathcal{P}_1, \mathcal{P}_2, \dots, \mathcal{P}_m\}$ of m convex d -polytopes lying on m parallel hyperplanes of $\mathbb{E}^{d+1}$, for any odd $d \geq 3$. Denoting by n_i the number of vertices of $\mathcal{P}_i$, we have shown that the worst-case complexity of $CH_{d+1}(\mathcal{P})$ is $O(\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor})$. This result suggests that, in order to compute $CH_{d+1}(\mathcal{P})$, it pays off to apply an output-sensitive convex hull algorithm to the set of vertices in $\mathcal{P}$. Indeed, for any odd $d \geq 5$ Seidel's shelling algorithm [27], or its modification by Matoušek and Schwarzkopf [21], results in a $O((\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor}) \log n)$ time algorithm, where $n = \sum_{i=1}^m n_i$. For $d = 3$, the divide-and-conquer algorithm by Chan, Snoeyink and Yap [11] can be competitive against Matoušek and Schwarzkopf's algorithm; hence, we may compute $CH_4(\mathcal{P})$ in $O(\min\{n^{4/3+\epsilon} + (\sum_{1 \leq i \neq j \leq m} n_i n_j) \log n, (\sum_{1 \leq i \neq j \leq m} n_i n_j) \log^2 n\})$ time, for any fixed $\epsilon > 0$. The above algorithms are nearly optimal for any odd $d \geq 3$; it remains an open problem to compute $CH_{d+1}(\mathcal{P})$ in worst-case optimal $O(\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor} + n \log n)$ time.

A direct consequence of our bound on the complexity of $CH_{d+1}(\mathcal{P})$ is a tight asymptotic bound on the worst-case complexity of the (weighted) Minkowski sum of two polytopes in any odd dimension $d \geq 3$. More precisely, consider a n -vertex d -polytope $\mathcal{P}$ and a m -vertex d -polytope $\mathcal{Q}$, and embed them on the hyperplanes $\{x_{d+1} = 0\}$ and $\{x_{d+1} = 1\}$ of $\mathbb{E}^{d+1}$, respectively. The weighted Minkowski sum $(1 - \lambda)\mathcal{P} \oplus \lambda\mathcal{Q}$, $\lambda \in (0, 1)$, is combinatorially equivalent to the intersection of $CH_{d+1}(\{\mathcal{P}, \mathcal{Q}\})$ with the hyperplane $\{x_{d+1} = \lambda\}$, whereas the Minkowski sum $\mathcal{P} \oplus \mathcal{Q}$ is nothing but $\frac{1}{2}\mathcal{P} \oplus \frac{1}{2}\mathcal{Q}$, scaled by a factor of 2. Applying our results, we deduce that the complexity of $(1 - \lambda)\mathcal{P} \oplus \lambda\mathcal{Q}$ (resp., $\mathcal{P} \oplus \mathcal{Q}$) is $\Theta(nm^{\lfloor \frac{d}{2} \rfloor} + mn^{\lfloor \frac{d}{2} \rfloor})$ for any odd $d \geq 3$. We would like to extend this tight bound to (weighted) Minkowski sums where the number of summands is greater than 2.

Capitalizing on our result on the complexity of convex hulls of convex polytopes lying on parallel hyperplanes, we have shown that the worst-case complexity of the convex hull $CH_d(\Sigma)$ of a set Σ of n spheres in $\mathbb{E}^d$ with a fixed number of m distinct radii $\rho_1, \rho_2, \dots, \rho_m$ is $O(\sum_{1 \leq i \neq j \leq m} n_i n_j^{\lfloor \frac{d}{2} \rfloor})$, for any odd $d \geq 3$, where n_i is the number of spheres with radius ρ_i . By means of an appropriate construction, described in Subsections 4.1 and 4.2, we have shown that the upper bound above is asymptotically tight, implying that our upper bound for $CH_{d+1}(\mathcal{P})$ is also tight. By slightly, but crucially, modifying the algorithm of Boissonnat *et al.* [6], $CH_d(\Sigma)$ may be computed in

$O(n^{\lfloor \frac{d}{2} \rfloor} + n \log n + T_{d+1}(n_1, \dots, n_m))$ time, where $T_{d+1}(n_1, \dots, n_m)$ stands for the time needed to compute the convex hull of m d -polytopes lying on m parallel hyperplanes in $\mathbb{E}^{d+1}$, where the i -th polytope has n_i vertices (cf. Section 5).

Finally, Boissonnat and Karavelas [7] have shown that convex hulls of spheres in $\mathbb{E}^d$ and additively weighted Voronoi cells in $\mathbb{E}^d$ are combinatorially equivalent. This equivalence suggests that we should be able to refine the worst-case complexity of an additively weighted Voronoi cell in any odd dimension, when the number of distinct radii of the spheres involved is considered fixed.

Acknowledgments

Partially supported by the FP7-REGPOT-2009-1 project “Archimedes Center for Modeling, Analysis and Computation”.

References

- [1] Nancy M. Amato and Edgar A. Ramos. On computing Voronoi diagrams by divide-prune-and-conquer. In *Proc. 12th Annu. ACM Sympos. Comput. Geom.*, pages 166–175, 1996.
- [2] A. M. Andrew. Another efficient algorithm for convex hulls in two dimensions. *Inform. Process. Lett.*, 9(5):216–219, 1979.
- [3] F. Aurenhammer. Power diagrams: properties, algorithms and applications. *SIAM J. Comput.*, 16(1):78–96, 1987.
- [4] David Avis, David Bremner, and Raimund Seidel. How good are convex hull algorithms? *Comput. Geom. Theory Appl.*, 7(5–6):265–302, 1997.
- [5] J.-D. Boissonnat, A. Cérézo, and J. Duquesne. An output-sensitive algorithm for constructing the convex hull of a set of spheres. In *IFIP Conference on Algorithms and efficient computation*, September 1992.
- [6] Jean-Daniel Boissonnat, André Cérézo, Olivier Devillers, Jacqueline Duquesne, and Mariette Yvinec. An algorithm for constructing the convex hull of a set of spheres in dimension d . *Comput. Geom. Theory Appl.*, 6(2):123–130, 1996.
- [7] Jean-Daniel Boissonnat and Menelaos I. Karavelas. On the combinatorial complexity of Euclidean Voronoi cells and convex hulls of d -dimensional spheres. In *Proc. 14th ACM-SIAM Sympos. Discrete Algorithms (SODA)*, pages 305–312, 2003.
- [8] Jean-Daniel Boissonnat and Mariette Yvinec. *Algorithmic Geometry*. Cambridge University Press, UK, 1998. Translated by Hervé Brönnimann.
- [9] T. M. Chan. Optimal output-sensitive convex hull algorithms in two and three dimensions. *Discrete Comput. Geom.*, 16(4):361–368, 1996.
- [10] T. M. Chan. Output-sensitive results on convex hulls, extreme points, and related problems. *Discrete Comput. Geom.*, 16(4):369–387, 1996.
- [11] T. M. Chan, J. Snoeyink, and Chee-Keng Yap. Primal dividing and dual pruning: Output-sensitive construction of four-dimensional polytopes and three-dimensional Voronoi diagrams. *Discrete Comput. Geom.*, 18(4):433–454, 1997.

- [12] D. R. Chand and S. S. Kapur. An algorithm for convex polytopes. *J. ACM*, 17(1):78–86, January 1970.
- [13] Bernard Chazelle. An optimal convex hull algorithm in any fixed dimension. *Discrete Comput. Geom.*, 10(1):377–409, 1993.
- [14] Bernard Chazelle and J. Matoušek. Derandomizing an output-sensitive convex hull algorithm in three dimensions. *Comput. Geom. Theory Appl.*, 5:27–32, 1995.
- [15] Jeff Erickson. New lower bounds for convex hull problems in odd dimensions. *SIAM J. Comput.*, 28(4):1198–1214, 1999.
- [16] L. R. Graham. An efficient algorithm for determining the convex hull of a finite planar set. *Inform. Process. Lett.*, 1(4):132–133, 1972.
- [17] Branko Grünbaum. *Convex Polytopes*, volume 221 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2003.
- [18] R. A. Jarvis. On the identification of the convex hull of a finite set of points in the plane. *Inform. Process. Lett.*, 2:18–21, 1973.
- [19] D. G. Kirkpatrick and R. Seidel. The ultimate planar convex hull algorithm? *SIAM J. Comput.*, 15:287–299, 1986.
- [20] Victor Klee. The number of vertices of a convex polytope. *Canadian J. Math.*, 16:701–720, 1964.
- [21] J. Matoušek and O. Schwarzkopf. Linear optimization queries. In *Proc. 8th Annu. ACM Sympos. Comput. Geom.*, pages 16–25, 1992.
- [22] P. McMullen. On a problem of Klee concerning convex polytopes. *Israel J. Math.*, 8(1):1–4, 1970.
- [23] Frank Nielsen and Mariette Yvinec. Output-sensitive convex hull algorithms of planar convex objects. *Internat. J. Comput. Geom. Appl.*, 8(1):39–66, 1998.
- [24] F. Preparata. An optimal real-time algorithm for planar convex hulls. *Commun. ACM*, 22:402–405, 1979.
- [25] F. P. Preparata and S. J. Hong. Convex hulls of finite sets of points in two and three dimensions. *Commun. ACM*, 20:87–93, 1977.
- [26] D. Rappaport. A convex hull algorithm for discs, and applications. *Comput. Geom. Theory Appl.*, 1(3):171–181, 1992.
- [27] Raimund Seidel. Constructing higher-dimensional convex hulls at logarithmic cost per face. In *Proc. 18th Annu. ACM Sympos. Theory Comput.*, pages 404–413, 1986.
- [28] Günter M. Ziegler. *Lectures on Polytopes*, volume 152 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.